arXiv:1007.5147v1 [math.CA] 29 Jul 2010

JACOB'S LADDERS AND THE NONLOCAL INTERACTION OF THE FUNCTION $Z^2(t)$ WITH THE FUNCTION $\tilde{Z}^2(t)$ ON THE DISTANCE $\sim (1-c)\pi(t)$ FOR THE COLLECTIONS OF DISCONNECTED SETS

JAN MOSER

Abstract. It is shown in this paper that there is a fine correlation of the fourth order between the functions $Z^2[\varphi_1(t)]$ and $\tilde{Z}^2(t)$, respectively. This correlation is with respect to two collections of disconnected sets. Corresponding new asymptotic formulae cannot be obtained within known theories of Balasubramanian, Heath-Brown and Ivic.

1. The result

In this paper we obtain some new properties of the signal

$$Z(t)=e^{i\vartheta(t)}\zeta\left(\frac{1}{2}+it\right)$$

generated by the Riemann zeta-function, where

$$\vartheta(t)=-\frac{t}{2}\ln\pi+\operatorname{Im}\ln\Gamma\left(\frac{1}{4}+i\frac{t}{2}\right)=\frac{t}{2}\ln\frac{t}{2\pi}-\frac{t}{2}-\frac{\pi}{8}+\mathcal{O}\left(\frac{1}{t}\right). \tag{1.1}$$

Let (see [3])

$$\begin{split} & G_3(x)=G_3(x;T,U)= \\ & =\bigcup_{T\leq g_{2\nu}\leq T+U}\{t:\ g_{2\nu}(-x)\leq t\leq g_{2\nu}(x)\},\ 0<x\leq\frac{\pi}{2}, \\ & G_4(y)=G_4(y;T,U)= \\ & =\bigcup_{T\leq g_{2\nu+1}\leq T+U}\{t:\ g_{2\nu+1}(-y)\leq t\leq g_{2\nu+1}(y)\},\ 0<y\leq\frac{\pi}{2}, \end{split} \tag{1.2}$$

$$U=T^{5/12}\ln^3 T, \tag{1.3}$$

and the collection of sequences $\{g_\nu(\tau)\}$, $\tau\in[-\pi,\pi]$, $\nu=1,2,\dots$ is defined by the equation (see [2], [3], (6))

$$\vartheta_1[g_\nu(\tau)]=\frac{\pi}{2}\nu+\frac{\tau}{2};\ g_\nu(0)=g_\nu,$$

where (comp. (1.1)

$$\vartheta_1(t)=\frac{t}{2}\ln\frac{t}{2\pi}-\frac{t}{2}-\frac{\pi}{8}.$$

Let

$$G_3(x)=\varphi_1[\mathring{G}_3(x)],\ G_4(y)=\varphi_1[\mathring{G}_4(y)], \tag{1.4}$$

Key words and phrases. Riemann zeta-function.

where $y = \varphi_1(T),\ T \geq T_0[\varphi_1]$ is the Jacob's ladder. The following theorem holds true.

Theorem.

$$
\begin{aligned}
&\int_{\overset{\circ}{G}_3(x)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t = \\
&\frac{x}{\pi}U\ln\frac{T}{2\pi} + \frac{2x}{\pi}\left(c + \frac{\sin x}{x}\right)U + \mathcal{O}(xT^{5/12}\ln^2 T), \\
&\int_{\overset{\circ}{G}_4(y)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t = \\
&\frac{y}{\pi}U\ln\frac{T}{2\pi} + \frac{2y}{\pi}\left(c - \frac{\sin y}{y}\right)U + \mathcal{O}(yT^{5/12}\ln^2 T),
\end{aligned}
\tag{1.5}
$$

where

$$
t - \varphi_1(t) \sim (1-c)\pi(t),\ t \to \infty,
\tag{1.6}
$$

and c is the Euler's constant and $\pi(t)$ is the prime-counting function.

Let (comp. (1.4)) $T = \varphi_1(\overset{\circ}{T}),\ T + U = \varphi_1(\overset{\circ}{\widehat{T+U}})$. Similarly to [14], (1.8) we have

$$
\rho\{[T, T+U]; [\overset{\circ}{T}, \overset{\circ}{\widehat{T+U}}]\} \sim (1-c)\pi(T),
\tag{1.7}
$$

where ρ stands for the distance of the corresponding segments.

Remark 1. Some nonlocal interaction of the functions $Z^2[\varphi_1(t)],\ \tilde{Z}^2(t)$ is expressed by eq. (1.5). This interaction is connected with two collections of disconnected sets unboundedly receding each from other (see (1.6), (1.7); $\rho \to \infty$ as $T \to \infty$) - like mutually receding galaxies (the Hubble law).

Remark 2. The asymptotic formulae (1.5) (comp. (1.3)) cannot be received by methods of Balasubramanian, Heath-Brown and Ivic (comp. [1]).

This paper is a continuation of the series of works [4]-[18].

2. ON BIG ASYMMETRY IN THE BEHAVIOUR OF THE FUNCTION $Z^2[\varphi_1(t)]\tilde{Z}^2(t)$ RELATIVELY TO THE SETS $\overset{\circ}{G}_3(x)$ AND $\overset{\circ}{G}_4(x)$

We obtain from (1.5)

Corollary.

$$
\begin{aligned}
&\int_{\overset{\circ}{G}_3(x)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t - \int_{\overset{\circ}{G}_4(x)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t = \\
&\frac{4}{\pi}U\sin x + \mathcal{O}(xT^{5/12}\ln^2 T),\ x \in (0, \pi/2],
\end{aligned}
\tag{2.1}
$$

especially, in the case $x = \pi/2$, we have

$$
\int_{\overset{\circ}{G}_3(\pi/2)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t - \int_{\overset{\circ}{G}_4(\pi/2)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t \sim \frac{4}{\pi}U,
\tag{2.2}
$$

where $[\overset{\circ}{T}, \overset{\circ}{\widehat{T+U}}] \subset \overset{\circ}{G}_3(\pi/2) \cup \overset{\circ}{G}_4(\pi/2)$.

Remark 3. The formulae (2.1), (2.2) represent the big difference of the areas (measures) of the figures which correspond to the functions

$$Z^2[\varphi_1(t)]\tilde{Z}^2(t),\ t\in\mathring{G}_3(x);\qquad Z^2[\varphi_1(t)]\tilde{Z}^2(t),\ t\in\mathring{G}_4(x).$$

The reason for this big asymmetry given by (2.2) is probably the fact that the zeroes of

$$\zeta\left(\frac{1}{2}+it\right),\ t\in\mathring{G}_3\left(\frac{\pi}{2}\right)\bigcup\mathring{G}_4\left(\frac{\pi}{2}\right)$$

lie preferably in the set $\mathring{G}_4(\pi/2)$.

3. Proof of the Theorem

3.1. The following lemma holds true (see [6], (2.5); [7], (3.3); [14], (4.1))

Lemma. For every integrable function (in the Lebesgue sense) $f(x),\ x\in[\varphi_1(T),\varphi_1(T+U)]$ the following is true

$$\int_T^{T+U} f[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{\varphi_1(T)}^{\varphi_1(T+U)} f(x)\mathrm{d}x,\ U\in\left(0,\frac{T}{\ln T}\right], \tag{3.1}$$

where $t-\varphi_1(t)\sim(1-c)\pi(t)$, c is the Euler's constant, $\pi(t)$ is the prime-counting function and

$$\tilde{Z}^2(t)=\frac{\mathrm{d}\varphi_1(t)}{\mathrm{d}t},\ \varphi_1(t)=\frac{1}{2}\varphi(t),$$
$$\tilde{Z}^2(t)=\frac{Z^2(t)}{2\Phi'_\varphi[\varphi(t)]}=\frac{Z^2(t)}{\left\{1+\mathcal{O}\left(\frac{\ln\ln t}{\ln t}\right)\right\}\ln t},$$

(see [7], (1.1), (3.1), (3.2)).

Remark 4. The formula (3.1) remains true also in the case when the integral on the right-hand side of (3.1) is only relatively convergent integral of the second kind (in the Riemann sense).

In the case (comp. (1.4) $T=\varphi_1(\mathring{T}),\ T+U=\varphi_1(\overset{\circ}{\widehat{T+U}})$) we obtain from (3.1)

$$\int_{\mathring{T}}^{\overset{\circ}{\widehat{T+U}}} f[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_T^{T+U} f(x)\mathrm{d}x. \tag{3.2}$$

3.2. First of all, we have from (3.2), for example

$$\int_{\mathring{g}_{2\nu(-x)}}^{\mathring{g}_{2\nu(x)}} f[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{g_{2\nu(-x)}}^{g_{2\nu(x)}} f(t)\mathrm{d}t \tag{3.3}$$

(see (1.4)). Next, in the case $f(t)=Z^2(t)$, we have the following $\tilde{Z}^2$-transformation

$$\begin{aligned}&\int_{\mathring{G}_3(x)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{G_3(x)} Z^2(t)\mathrm{d}t,\\&\int_{\mathring{G}_4(y)} Z^2[\varphi_1(t)]\tilde{Z}^2(t)\mathrm{d}t=\int_{G_4(y)} Z^2(t)\mathrm{d}t.\end{aligned} \tag{3.4}$$

Let us remind that we have proved in the paper [3] the following mean-value formulae

$$\begin{aligned} & \int_{G_3(x)} Z^2(t)\mathrm{d}t=\frac{x}{\pi}U\ln\frac{T}{2\pi}+\frac{2x}{\pi}\left(c+\frac{\sin x}{x}\right)U+\mathcal{O}(xT^{5/12}\ln^2T), \\ & \int_{G_4(y)} Z^2(t)\mathrm{d}t=\frac{y}{\pi}U\ln\frac{T}{2\pi}+\frac{2y}{\pi}\left(c-\frac{\sin y}{y}\right)U+\mathcal{O}(yT^{5/12}\ln^2T). \end{aligned} \tag{3.5}$$

Now, the formula (1.5) follows from (3.4), (3.5).

Remark 5. The formulae (3.5) are the consequences of their discrete form

$$\begin{aligned} & \sum_{T\leq g_{2\nu}\leq T+U} Z^2[g_{2\nu}(\tau)]= \\ & \frac{1}{2\pi}U\ln^2\frac{T}{2\pi}+\frac{1}{\pi}(c+\cos\tau)U\ln\frac{T}{2\pi}+\mathcal{O}(T^{5/12}\ln^3T), \\ & \sum_{T\leq g_{2\nu+1}\leq T+U} Z^2[g_{2\nu+1}(\tau)]= \\ & \frac{1}{2\pi}U\ln^2\frac{T}{2\pi}+\frac{1}{\pi}(c-\cos\tau)U\ln\frac{T}{2\pi}+\mathcal{O}(T^{5/12}\ln^3T) \end{aligned}$$

(see [3], (10)).

I would like to thank Michal Demetrian for helping me with the electronic version of this work.

References

[1] A. Ivic, 'The Riemann zeta-function', A Willey-Interscience Pub., New York, 1985.

[2] J. Moser, 'An improvement of the theorem of Hardy-Littlewood on density of zeroes of the function $\zeta\left(\frac{1}{2}+it\right)$', Acta Arithmetica, 43 (1983), 21-47 (in russian).

[3] J. Moser, 'New mean-value theorems for the function $|\zeta\left(\frac{1}{2}+it\right)|^2$', Acta Math. Univ. Comen., 46-47 (1985), pp. 21-40, (in russian).

[4] J. Moser, 'Jacob's ladders and the almost exact asymptotic representation of the Hardy-Littlewood integral', (2008), arXiv:0901.3973.

[5] J. Moser, 'Jacob's ladders and the tangent law for short parts of the Hardy-Littlewood integral', (2009), arXiv:0906.0659.

[6] J. Moser, 'Jacob's ladders and the multiplicative asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral', (2009), arXiv:0907.0301.

[7] J. Moser, 'Jacob's ladders and the quantization of the Hardy-Littlewood integral', (2009), arXiv:0909.3928.

[8] J. Moser, 'Jacob's ladders and the first asymptotic formula for the expression of the sixth order $|\zeta(1/2+i\varphi(t)/2)|^4|\zeta(1/2+it)|^2$', (2009), arXiv:0911.1246.

[9] J. Moser, 'Jacob's ladders and the first asymptotic formula for the expression of the fifth order $Z[\varphi(t)/2+\rho_1]Z[\varphi(t)/2+\rho_2]Z[\varphi(t)/2+\rho_3]\hat{Z}^2(t)$ for the collection of disconnected sets', (2009), arXiv:0912.0130.

[10] J. Moser, 'Jacob's ladders, the iterations of Jacob's ladder $\varphi_1^k(t)$ and asymptotic formulae for the integrals of the products $Z^2[\varphi_1^n(t)]Z^2[\varphi^{n-1}(t)]\cdots Z^2[\varphi_1^0(t)]$ for arbitrary fixed $n\in\mathbb{N}$' (2010), arXiv:1001.1632.

[11] J. Moser, 'Jacob's ladders and the asymptotic formula for the integral of the eight order expression $|\zeta(1/2+i\varphi_2(t))|^4|\zeta(1/2+it)|^4$', (2010), arXiv:1001.2114.

[12] J. Moser, 'Jacob's ladders and the asymptotically approximate solutions of a nonlinear diophantine equation', (2010), arXiv: 1001.3019.

[13] J. Moser, 'Jacob's ladders and the asymptotic formula for short and microscopic parts of the Hardy-Littlewood integral of the function $|\zeta(1/2+it)|^4$', (2010), arXiv:1001.4007.

[14] J. Moser, 'Jacob's ladders and the nonlocal interaction of the function $|\zeta(1/2+it)|$ with $\arg\zeta(1/2+it)$ on the distance $\sim(1-c)\pi(t)$', (2010), arXiv: 1004.0169.
[15] J. Moser, 'Jacob's ladders and the $\tilde{Z}^2$ - transformation of polynomials in $\ln\varphi_1(t)$', (2010), arXiv: 1005.2052.
[16] J. Moser, 'Jacob's ladders and the oscillations of the function $|\zeta\left(\frac{1}{2}+it\right)|^2$ around the main part of its mean-value; law of the almost exact equality of the corresponding areas', (2010), arXiv: 1006.4316
[17] J. Moser, 'Jacob's ladders and the nonlocal interaction of the function $Z(t)$ with the function $\tilde{Z}^2(t)$ on the distance $\sim(1-c)\pi(t)$ for a collection of disconneted sets', (2010), arXiv: 1006.5158
[18] J. Moser, 'Jacob's ladders and the $\tilde{Z}^2$-transformation of the orthogonal system of trigonometric functions', (2010), arXiv: 1007.0108.

DEPARTMENT OF MATHEMATICAL ANALYSIS AND NUMERICAL MATHEMATICS, COMENIUS UNIVERSITY, MLYNSKA DOLINA M105, 842 48 BRATISLAVA, SLOVAKIA
E-mail address: `jan.mozer@fmph.uniba.sk`